


# SPOZNAJTE OPIČIE KIAHNE



ÚRAD  
VEREJNÉHO  
ZDRAVOTNÍCTVA  
SLOVENSKEJ  
REPUBLIKY

**Opičie kiahne sa môžu prenášať blízky fyzickým kontaktom (koža na kožu), a to aj prostredníctvom intímneho kontaktu alebo počas sexuálneho styku.**



## AKO SA OCHORENIE ŠÍRI



Úzkym kontaktom s:



**vyrážkami ako aj výlučkami z nich**



**kontaminovanými predmetmi**

ide o predmety ako napr. uteráky, oblečenie, riady, posteľná bielizeň, sexuálne hračky




**telesnými tekutinami alebo veľkými kvapôčkami priamo z tváre do tváre**


je potrebný dlhší kontakt než v prípade COVID-19

**!** Inkubačný čas môže byť 5 až 21 dní

## AKÉ SÚ PRÍZNAKY OCHORENIA




**Horúčka alebo zimnica**



**Opuch lymfatických uzlín**



**Silná bolesť hlavy**



**Bolesti svalov či chrbta**



**Únava alebo vyčerpanie**



**Vyrážka sa objaví zvyčajne 1 až 3 dni od výskytu niektorého z príznakov**

Pľuzgieriky sa môžu objaviť aj na dlaniach, chodidlách, vnútri úst, pri análnom otvore alebo na genitáliách. Môže ísť aj o ojedinelý pľuzgierik.

**Infekcia sa môže ďalej prenášať až do zahojenia poslednej chrasty** **!**


**Pri podozrení na ochorenie alebo úzky kontakt s chorým konzultujte vhodný postup s lekárom. Ak je potrebné vyšetrenie, vyrážku si mimo domácnosti prekryte oblečením a nasadte si respirátor.**

# RECOGNIZING MONKEYPOX



PUBLIC  
HEALTH  
AUTHORITY  
OF THE SLOVAK  
REPUBLIC

**Monkeypox can be transmitted through close physical contact (skin-to-skin), including intimate or sexual contact.**



## HOW THE DISEASE SPREADS

Close contact with:




**!** The incubation period ranges from 5 to 21 days



**contaminated objects**

items such as towels, clothes, dishes, bed linen, sex toys, etc.



**rash, including secretions from pox lesions**


**body fluids or large droplets during close (face-to-face) contact**

longer contact is needed compared to COVID-19

## WHAT ARE THE SYMPTOMS




**Fever or chills**



**Swollen lymph nodes**



**Strong headache**



**Muscle aches or backache**



**Fatigue or exhaustion**

**The rash (pimples or blisters) typically appears 1 to 3 days after initial symptoms.**



Blisters can also appear on the palms of the hands, feet soles, inside the mouth, near the anus or on the genitals. It can also be a single blister.

**!** Infection can be transmitted until the last scab heals

**If you have symptoms or if you had close contact with the infected person, you should talk to your healthcare provider. If personal examination is necessary, cover the rash with clothing and wear a respirator when leaving home.**